

**IN THE SPECIFICATION**

Please amend the title as follows:

~~NOVEL~~ POLYPEPTIDES INVOLVED IN IMMUNE RESPONSE

Please replace the Abstract with the replacement Abstract included, herewith.

Please cancel the Amendment dated October 28, 2002.

Please replace the paragraph beginning at page 5, line 4, and ending at page 5, line 11, with the following:

b) the nucleotide sequence encoding the polypeptide as set forth in Figure 2A (SEQ ID NO: 6) from residues 1-322 or from residues 47-322 or as set forth in Figure 3A (SEQ ID NO: 11) from residues 1-288 or from residues 19-288, 20-288, 21-288, 22-288, 24-288, or 28-288; or as set forth in Figure 12A (SEQ ID NO: 16) from residues 1-302 or from residues 19-302, 20-302, 21-302, 22-302, 24-302 or 28-302;

Please replace the paragraph beginning at page 20, line 19, and ending at page 20, line 36, with the following:

CD28 related protein-1, or CRP1, is predicted to be a type I transmembrane protein with a signal sequence and extracellular domain at the amino-terminus, a transmembrane domain, and a carboxy terminal intracellular domain (Figure 1). The full-length CRP1 protein is 180 amino acids in its mature form. The predicted leader sequence spans about amino acid residues 1-20 (relative to the initiating methionine) and the extracellular domain of the mature protein encompasses about residues 21-145

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com

E<sub>2</sub>

(Example 1). The predicted transmembrane domain spans about residues 146-163 and the intracellular domain encompasses about residues 164-200. The amino terminal extracellular domain is similar to an Ig loop with conserved putative intra- and inter-molecular bonding cysteines. Furthermore, a "MYPPPY" motif (SEQ ID NO: 36), which is previously known to be important for B7.1 and B7.2 binding to CD28 and CTLA-4, is also partially conserved.

Please replace the paragraph beginning at page 21, line 13, and ending at page 21, line 27, with the following:

E<sub>3</sub>

Human CRP1 is a transmembrane protein having the nucleotide and amino acid sequence as shown in Figure 13A (SEQ ID NOs: 21 and 22, respectively). The predicted leader sequence spans about residues 1-19 or about residues 1-20. The predicted mature amino terminus is at residues 20 or 21. Preferably, the mature amino terminus is at position 21. The extracellular domain spans from any of the predicted mature amino termini to about amino acid residue 140, the transmembrane domain spans about residues 141-161 and the intracellular domain spans about residues 162-199. Human CRP1 protein has 69% identity to the murine protein and the corresponding nucleotide sequences are 77% identical. The sequence of human CRP1 was reported in Hutloff et al. Nature 397, 263-266 (1999).

Please replace the paragraph beginning at page 40, line 14, and ending at page 40, line 26, with the following:

E  
4

The term "CRP1 or B7RP1 polypeptide" refers to a polypeptide having the amino acid sequence of Figure 1A (SEQ ID NO:2), Figure 2A (SEQ ID NO:7) or Figure 3A (SEQ ID NO:12), Figure 12A (SEQ ID NO: 17), or Figure 13A (SEQ ID NO: 22), and all related polypeptides described herein. Related polypeptides includes allelic variants, splice variants, fragments, derivatives, substitution, deletion, and insertion variants, fusion polypeptides, and orthologs. Such related polypeptides may be mature polypeptides, *i.e.*, polypeptide lacking a signal peptide. A CRP1 or B7RP1 polypeptide may or may not have amino terminal methionine, depending on the manner in which they are prepared.

Please replace the paragraph beginning at page 40, line 27, and ending at page 41, line 8, with the following:

E  
5

The term "CRP1 or B7RP1 polypeptide fragment" refers to a peptide or polypeptide that is less than the full length amino acid sequence of a CRP1 or B7RP1 polypeptide as set forth in Figure 1A (SEQ ID NO:2), Figure 2A (SEQ ID NO:7) or Figure 3A (SEQ ID NO:12), Figure 12A (SEQ ID NO: 17), or Figure 13A (SEQ ID NO: 22). Such a fragment may result from truncation at the amino terminus, truncation at the carboxy terminus, and/or a deletion internal to the polypeptide sequence. Such CRP1 or B7RP1 polypeptides fragments may be prepared with or without an amino terminal methionine. In addition, CRP1 or B7RP1 polypeptides fragments may be naturally-occurring splice variants, other splice variants, and fragments resulting from naturally occurring *in vivo* protease activity. Preferred CRP1 or B7RP1 polypeptide fragments include soluble forms of CRP1 or B7RP1 which lack a functional

E5  
Cont

transmembrane domain and comprise part or all of the extracellular domain of either CRP1 or B7RP1.

Please replace the paragraph beginning at page 41, line 9, and ending at page 41, line 20, with the following:

E6

The term "CRP1 or B7RP1 polypeptide variants" refers to CRP1 or B7RP1 polypeptides whose amino acid sequences contain one or more amino acid sequence substitutions, deletions, and/or additions as compared to the CRP1 or B7RP1 polypeptides amino acid sequences set forth in Figure 1A (SEQ ID NO:2), Figure 2A (SEQ ID NO:7) or Figure 3A (SEQ ID NO:12), Figure 12A (SEQ ID NO: 17), or Figure 13A (SEQ ID NO: 22). Such CRP1 or B7RP1 polypeptides variants can be prepared from the corresponding CRP1 and B7RP1 polypeptides nucleic acid molecule variants, which have a DNA sequence that varies accordingly from the DNA sequences for CRP1 or B7RP1 polypeptides.

Please replace the paragraph beginning at page 92, line 28, and ending at page 93, line 14, with the following:

E7

A cDNA clone containing an open reading frame of 199 amino acids was obtained (Figure 13A). This cDNA clone contained nucleotide and amino acid homologies to the murine CRP1 clone described in Example 1 and Figure 1. The nucleotides corresponding to the open reading frame of this human clone was 77% identical to the murine CRP1 gene. Translation of the human sequence and subsequent comparison with the murine CRP1 protein revealed 69% amino acid identity with the murine protein (Figure 13B). In addition, the motif between amino acids 114 to

E7  
Cont

119, "FDPPPF" (SEQ ID NO: 37), was conserved between the murine and human CRP1 genes. This motif corresponds to the "MYPPPY" (SEQ ID NO: 36) motif in murine and human CD28 that is essential for B7 protein interaction. Furthermore, the cysteines at amino acid positions 42, 109, and 141 are also conserved. These cysteines correspond to cysteines in CD28 and CTLA-4 that are involved in Ig loop formation and intermolecular disulfide dimerization. The close similarity with murine CRP1, and structural similarities with the CD28 homology family, indicate that this is the human CRP1 homolog.

Please amend the paragraphs beginning on page 9, line <sup>34</sup>4 and ending on page 10, line 24 as follows:

E8

Figures 1—A.1-1A.2) (SEQ ID NOS: 1&2) DNA and amino acid sequence of murine CRP1 (mCRP1). Predicted signal sequence of CRP1 is underlined at the amino-terminus and the experimentally determined pro-peptide cleavage site is indicated by an asterisk. Predicted transmembrane sequence is underlined toward the carboxy-terminus. Figure 1B) (SEQ ID NOS: 3,4, 5) Amino acid alignment of B7RP1 protein sequence (mCRP1) with murine CD28.

Figures 2—A.1-2A.2) (SEQ ID NOS: 6 & 7) DNA and amino acid sequence of murine B7RP1 (mB7RP1). Predicted signal sequence of B7RP1 is underlined at the amino-terminus and the experimentally determined pro-peptide cleavage site is indicated by an asterisk. Predicted transmembrane sequence is underlined toward the carboxy-terminus. Figure 2B) (SEQ ID NOS: 8, 9 & 10) Amino acid alignment of B7RP1 protein sequence (mB7RP1) with murine CD80 (mCD80).

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com

E<sub>8</sub>

Figures 3.—A.1-3A.2) (SEQ ID NOS: 11 & 12) Structure and sequence of the protein coding region of the putative human B7RP1 (hB7RP1). Predicted signal sequence of hB7RP1 is underlined at the amino-terminus. Predicted signal peptide cleavage sites are marked by asterisks. Predicted transmembrane sequence is underlined toward the carboxy-terminus. Figure 3B) (SEQ ID NOS: 13, 14 & 15) Amino acid alignment of the putative mature hB7RP1 protein with the mature murine B7RP1 (mB7RP1) protein.

✓  
Please amend the paragraphs beginning on page 14, line 32 and ending on page 15, line 12 as follows:

E<sub>9</sub>

Figures 12.—A.1-12A.2) (SEQ ID NOS: 16 & 17) Structure and sequence of the protein coding region of human B7RP1 (hB7RP1). Predicted signal sequence of hB7RP1 is underlined at the amino-terminus. Predicted signal peptide cleavage sites are marked by asterisks. Predicted transmembrane sequence is underlined toward the carboxy-terminus. Figure 12B) (SEQ ID NOS: 18, 19 & 20) Amino acid alignment of the putative mature hB7RP1 protein (human) with the mature murine B7RP1 (mB7RP1) protein (mouse).

Figures 13.—A.1-13A.2) (SEQ ID NOS: 21 & 22) Structure and sequence of the protein coding region of human CRP1 (hCRP1). Predicted signal sequence of hCRP1 is underlined at the amino-terminus. Predicted signal peptide cleavage sites are marked by asterisks. Predicted transmembrane sequence is underlined toward the carboxy-terminus. Figure 13B) (SEQ ID NOS: 23 & 24) Amino acid alignment of the hCRP1 protein with the murine CRP1 (~~mB7RP1~~mCRP1) protein.

FINNEGAN  
HENDERSON  
FARABOW  
GARRETT &  
DUNNER LLP

1300 I Street, NW  
Washington, DC 20005  
202.408.4000  
Fax 202.408.4400  
www.finnegan.com